


SURFICIAL GEOLOGY OF THE BAINBRIDGE QUADRANGLE, OHIO

by
Richard R. Pavey
with GIS and Cartography by Dean R. Martin



- UNIT DESCRIPTIONS
- Water. Large modern lakes and reservoirs.
 - Made land. Large cut-and-fill areas.
 - Sand-and-gravel pit.
 - Quarry.
 - a Alluvium (Holocene). Found within floodplains of modern streams.
 - c Thick colluvium and landslide debris (Holocene and Pleistocene?).
 - L Clay and silt (Wisconsinan). Massive to laminated; may contain thin, interbedded, fine sand and minor gravel. Deposited in lakebeds at the ice margin.
 - SG Sand and gravel (Wisconsinan). Stratified sand and gravel with thin, discontinuous layers of silt and clay. Deposited as valley trains and outwash plains.
 - Cl Clay and silt (Illinoian). Massive to laminated; may contain thin, interbedded, fine sand and minor gravel; deeply weathered, leached, and dissected. Deposited in lakebeds at the ice margin.
 - SGi Sand and gravel (Illinoian). Similar to SG above, but more deeply weathered, leached, and dissected.
 - ICI Ice-contact deposits (Illinoian). Highly variable deposits of poorly sorted gravel and sand; silt, clay, and till lenses common; may be partially covered or surrounded by till. Deposited directly from stagnant ice as kame or esker landforms; deeply weathered, leached, and dissected.
 - Ti Loam till (Illinoian). Deeply weathered, leached, and dissected. Overlain by up to 4 feet of loess where not eroded. Till may contain silt, sand, and gravel lenses. Deposited directly from ice.
 - Mc Cuyahoga Formation (Mississippian). Sandstone, siltstone, and shale; thin to massive bedding; mainly shades of gray, olive, brown, and yellow.
 - Db Berea Sandstone (Devonian). Gray to shades of brown; medium grained to silty.
 - Dbd Bedford Shale (Devonian). Gray, red, and brown; silty to clayey; locally abundant siltstone and sandstone interbeds.
 - Doh Ohio Shale (Devonian). Mostly black; carbonaceous; calcareous concretions common in lower portion.
 - Stb Tymochtee, Greenfield, and Peebles Dolomites and Lilley and Bisher Formations undivided (Silurian). Thin to thick, interbedded dolomites and shales.
- Significant landslide



Location of Bainbridge 1:24,000 quadrangle in Ohio.

The products of the Ohio Department of Natural Resources (ODNR), Division of Geological Survey, including digital maps and printed maps and any other associated documents, are intended to provide general geologic information only and should not be used for any other purpose. They are not intended for resale or to replace site-specific investigations. These data were compiled by ODNR Division of Geological Survey, which reserves the publication rights to this material. If these data are used in the compilation of other data sets or maps for distribution or publication, this source must be referenced.

Neither the Ohio Department of Natural Resources, nor any agency thereof, nor any of their employees, contractors, or subcontractors, make any warranty, express or implied, nor assume any legal liability or responsibility for the accuracy, completeness, or usefulness of this product. Any use thereof for a purpose other than for which said information or product was intended shall be solely at the risk of the user.



www.OhioGeology.com



Basemap derived from various State of Ohio datasets
Projection is Ohio coordinate system, south zone
North American Datum 1983

